

**This Bluebird Label is Intended for Use Only with Monensin¹, Tylosin²
and Ractopamine³**

Liquid Medicated Heifer Supplement

Type C Medicated Feed For Beef and Dairy Heifers

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

INDICATIONS

Heifers Fed in Confinement for Slaughter: For increased Rate of Weight Gain, Improved Feed Efficiency and Suppression of Estrus (Heat).

ACTIVE DRUG INGREDIENTS

Melengestrol acetate⁴ (0.125-1.0) mg /lb **

GUARANTEED ANALYSIS

Crude Protein, not less than	_____	%
Non-Protein Nitrogen (NPN) ⁵ , not more than	_____	%
Crude Fat, not less than	_____	%
Crude Fiber, not more than	_____	%
Calcium, not less than	_____	%
Calcium, not more than	_____	%
Phosphorus, not less than.....	_____	%
Salt ⁶ , not less than	_____	%
Salt ⁶ , not more than.....	_____	%
Sodium ⁷ , not less than	_____	%
Sodium ⁷ , not more than.....	_____	%
Potassium, not less than	_____	%
Vitamin A ^{6,8} , not less than.....	_____	I.U./lb
pH.....	4.0 to 8.0	

⁵When added.

⁶If added.

⁷Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

⁸Other than precursors of Vitamin A.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

DIRECTIONS FOR USE

Must be top dressed or mixed with a complete ration containing monensin (10 to 40 g/ton), tylosin (8 to 10 g/ton) and ractopamine (9.8 to 24.6 g/ton).

Feed at the rate of 0.5-2.0 pound(s) per head per day (specify one level) to provide 0.25-0.5 mg melengestrol acetate per head per day (specify one level). Feed melengestrol acetate in this combination for the final 28 to 42 days on feed.

MIXING DIRECTIONS

Mixing directions for liquid Type C feeds stored in recirculation tank systems are: "Recirculate immediately prior to use for no less than 10 minutes, moving not less than 1 percent of the tank contents from the bottom of the tank to the top. Recirculate daily, as directed in this paragraph even when the Type C feed is not used." Mixing directions for liquid Type C feeds stored in mechanical, air or other agitation- type tank systems are: "Agitate immediately prior to use for not less than 10 minutes, creating a turbulence at the bottom of the tank that is visible at the top. Agitate daily, as directed in this paragraph, even when the Type C feed is not used."



CAUTION

Melengestrol Acetate is not effective in steers and spayed heifers.

Withdrawal periods of three to five days should be avoided to prevent the possibility that the heifers may come into estrus (heat) at the time of loading.

Do not use in any finished feed (supplement, concentrate or complete feed) containing in excess of 2% bentonite.

WARNING

 A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal. 

Approved by FDA under ANADA # 200-649

MANUFACTURED BY
BLUE BIRD FEED MILL
Any town, USA 12345

Net Weight lb (kg) on bag or bulk

Lot No. _____

**The medicated feed label must state a single drug concentration.

¹Sourced from Monovet[®], ANADA # 200-639

²Sourced from Tylovet[®], ANADA # 200-484

³Sourced from Actogain[®], ANADA # 200-548

⁴Sourced from MGA[®], NADA # 039-402

Monovet and Tylovet are registered trademarks of Huvepharma EOOD.
Actogain and MGA are registered trademarks of Zoetis Inc.